

On quantizing semisimple basic algebras, II: The general case

Mark J. Gotay*

Department of Mathematics
University of Hawai'i
2565 The Mall
Honolulu, HI 96822 USA
E-mail: gotay@math.hawaii.edu

November 30, 2001
(Revised February 7, 2008)

Abstract

We prove that there is no consistent polynomial quantization of the coordinate ring of a basic non-nilpotent coadjoint orbit of a semisimple Lie group.

*Supported in part by NSF grant DMS 00-72434.

1 Introduction

In a recent paper (Gotay [2002]) we showed that there do not exist polynomial quantizations of the coordinate ring $P(M)$ of a basic semisimple coadjoint orbit $M \subset \mathfrak{sl}(2, \mathbb{R})^*$. Here we extend this result to any basic non-nilpotent coadjoint orbit of a general semisimple Lie group:

Theorem 1.1 *Let $\mathfrak{b}$ be a finite-dimensional semisimple Lie algebra, and M a basic non-nilpotent coadjoint orbit in $\mathfrak{b}^*$. Then there are no polynomial quantizations of the coordinate ring $P(M)$.*

We refer the reader to Gotay [2000] for definitions and discussions of basic coadjoint orbits and quantization.

In particular, a coadjoint orbit $M \subset \mathfrak{b}^*$ is *basic* provided $\mathfrak{b}$ has no subalgebras which act transitively on M and which globally separate points of M . Unfortunately, it is difficult to determine exactly which orbits are basic. From §4 in Gotay, Grabowski, and Grundling [2000] and Prop. 2.1 in Gotay [2002] we know that $M \subset \mathfrak{b}^*$ will be basic for a semisimple Lie algebra $\mathfrak{b}$ whenever

- (i) $\mathfrak{b}$ is compact and M is principal,
- (ii) $\mathfrak{b}$ is compact and simple, and
- (iii) M is principal nilpotent.

Consider the symmetric algebra $S(\mathfrak{b})$, regarded as the ring of polynomials on $\mathfrak{b}^*$. The Lie bracket on $\mathfrak{b}$ may be extended via the Leibniz rule to a Poisson bracket on $S(\mathfrak{b})$, so that the latter becomes a Poisson algebra. Let $I(M)$ be the associative ideal in $S(\mathfrak{b})$ consisting of all polynomials which vanish on M and set $P(M) = S(\mathfrak{b})/I(M)$. Since M is an orbit $I(M)$ is also a Lie ideal, hence a Poisson ideal, so the coordinate ring $P(M)$ of M inherits the structure of a Poisson algebra from $S(\mathfrak{b})$. We denote the Poisson brackets on both $P(M)$ and $S(\mathfrak{b})$ by $\{\cdot, \cdot\}$.

Here we are interested in quantizing the coordinate ring $P(M)$. By a *quantization* of $P(M)$ we mean a Lie representation $\mathcal{Q}$ thereof by symmetric operators preserving a fixed dense domain D in some separable Hilbert space $\mathcal{H}$, such that $\mathcal{Q} \upharpoonright \mathfrak{b}$ is irreducible, integrable, and faithful. Let $\mathcal{A}$ be the associative operator algebra generated over $\mathbb{C}$ by I and $\{\mathcal{Q}(b) \mid b \in \mathfrak{b}\}$. We say that a quantization $\mathcal{Q}$ of $P(M)$ is *polynomial* if $\mathcal{Q}$ is valued in $\mathcal{A}$.

2 Proof of Theorem 1.1

Suppose to the contrary that $\mathcal{Q}$ were a polynomial quantization of $P(M)$ in a dense invariant domain D in a Hilbert space $\mathcal{H}$. By extending $\mathcal{Q}$ to be complex linear, we obtain a Lie representation $\mathcal{Q}_{\mathbb{C}}$ of the Poisson algebra $P(M, \mathbb{C})$ of complex-valued polynomials on M in D .

By assumption the representation of $\mathfrak{b}$ in D provided by $\mathcal{Q}$ may be integrated to a strongly continuous unitary representation Π of the 1-connected Lie group B with Lie algebra $\mathfrak{b}$ in $\mathcal{H}$. Let $B_{\mathbb{C}}$ be the universal complexification of B ; since B is simply connected, $B_{\mathbb{C}}$ can be identified with the 1-connected semisimple complex analytic group with Lie algebra the complexification $\mathfrak{b}_{\mathbb{C}}$ of $\mathfrak{b}$. (See Varadarajan [1984], pps. 256–258 and 400–404 for background on complexifications of Lie groups.) Since B is semisimple, B is a closed subgroup of $B_{\mathbb{C}}$, and so we may use induction to obtain a strongly continuous unitary representation $\Pi_{\mathbb{C}}$ of $B_{\mathbb{C}}$ in a certain infinite-dimensional Hilbert space $\mathcal{K}$.

Now let C be a compact real form of $B_{\mathbb{C}}$, and denote by Γ the restriction of $\Pi_{\mathbb{C}}$ to C . As every strongly continuous unitary representation of a compact Lie group is completely reducible, we may decompose

$$\mathcal{K} = \bigwedge_{i \in I} \bigoplus \mathcal{K}_i$$

for some index set $I \subset \mathbb{Z}$, where the finite-dimensional invariant subspaces $\mathcal{K}_i$ are the carriers of the irreducible constituents Γ_i of Γ . Let $\mathfrak{c}$ be the Lie algebra of C ; then for each $i \in I$, we have the derived representation $d\Gamma_i$ of $\mathfrak{c}$ in $\mathcal{K}_i$. Set $d\Gamma = \bigoplus_{i \in I} d\Gamma_i$; this gives a representation of $\mathfrak{c}$ in the dense subspace

$$D_C = \bigoplus_{i \in I} \mathcal{K}_i.$$

Choose a basis $\{c_1, \dots, c_r\}$ of $\mathfrak{c}$. Since $\mathfrak{c}_{\mathbb{C}} = \mathfrak{b}_{\mathbb{C}}$ and as by assumption $\mathcal{Q}$ is valued in $\mathcal{A}$, for every $f \in P(M, \mathbb{C})$ we may expand

$$\mathcal{Q}_{\mathbb{C}}(f) = \sum_{n_1, \dots, n_r} a_{n_1, \dots, n_r}^f \mathcal{Q}_{\mathbb{C}}(c_1)^{n_1} \cdots \mathcal{Q}_{\mathbb{C}}(c_r)^{n_r}$$

for some coefficients $a_{n_1, \dots, n_r}^f$. By means of this formula we can extend the representation $d\Gamma$ of $\mathfrak{c}$ to a Lie representation γ of $P(M, \mathbb{C})$ in D_C :

$$\gamma(f) = \sum_{n_1, \dots, n_r} a_{n_1, \dots, n_r}^f d\Gamma(c_1)^{n_1} \cdots d\Gamma(c_r)^{n_r}$$

with the *same* coefficients. As each subspace $\mathcal{K}_i$ is invariant, γ restricts to a representation γ_i of $P(M, \mathbb{C})$ in $\mathcal{K}_i$. We will show that the existence of these representations γ_i leads to a contradiction.

To this end we recall the following algebraic fact, the proof of which is given in Gotay, Grabowski, and Grundling [2000].

Lemma 2.1 *If L is a finite-codimensional Lie ideal of an infinite-dimensional Poisson algebra P with identity, then either L contains the derived ideal $\{P, P\}$ or there is a maximal finite-codimensional associative ideal J of P such that $\{P, P\} \subset J$.*

We apply Lemma 2.1 to each $L_i = \ker \gamma_i$ which, as $\mathcal{K}_i$ is finite-dimensional, has finite codimension in $P = P(M, \mathbb{C})$. First suppose there is an i for which $\{P, P\} \not\subset L_i$. Then there must exist a maximal finite-codimensional associative ideal J_i in P with $\{P, P\} \subset J_i$. If ρ is the projection $S(\mathfrak{b}_{\mathbb{C}}) \rightarrow P$, then $I_i = \rho^{-1}(J_i)$ is a maximal finite-codimensional associative ideal in $S(\mathfrak{b}_{\mathbb{C}})$ with $\{S(\mathfrak{b}_{\mathbb{C}}), S(\mathfrak{b}_{\mathbb{C}})\} \subset I_i$. Since by semisimplicity

$$\mathfrak{b}_{\mathbb{C}} = \{\mathfrak{b}_{\mathbb{C}}, \mathfrak{b}_{\mathbb{C}}\} \subset \{S(\mathfrak{b}_{\mathbb{C}}), S(\mathfrak{b}_{\mathbb{C}})\} \subset I_i,$$

and since $1 \notin I_i$ (as I_i is proper), it follows that I_i is the associative ideal generated by $\mathfrak{b}_{\mathbb{C}}$. (Actually, this shows that $S(\mathfrak{b}_{\mathbb{C}}) = \mathbb{C} \oplus I_i$.)

Since the orbit M is not nilpotent, there is a nonzero Casimir $\Omega \in S(\mathfrak{b}_{\mathbb{C}})$, i.e. $\rho(\Omega) = \omega$ for some constant $\omega \neq 0$. Since $\mathfrak{b}_{\mathbb{C}}$ is semisimple it follows from the above observations that $\Omega \in I_i$. But then $\Omega - \omega \notin I_i$, which is a contradiction since $\Omega - \omega \in \ker \rho \subset I_i$.

Thus for *every* i it must be the case that $\{P, P\} \subset L_i$. Again semisimplicity gives $\mathfrak{b}_{\mathbb{C}} = \{\mathfrak{b}_{\mathbb{C}}, \mathfrak{b}_{\mathbb{C}}\} \subset L_i$, and so $\gamma \upharpoonright \mathfrak{b}_{\mathbb{C}} = 0$. In particular, then, $d\Gamma = 0$. Since $\mathfrak{c}$ is a compact real form of $\mathfrak{b}_{\mathbb{C}}$, the Cartan decomposition of $\mathfrak{b}_{\mathbb{C}}$ implies that $d\Pi_{\mathbb{C}} = 0$. It follows from the induction construction that the original derived representation $d\Pi$ of $\mathfrak{b}$ in the domain D must be zero as well. But then $\mathcal{Q} \upharpoonright \mathfrak{b} = 0$, which contradicts the requirement that a quantization represent $\mathfrak{b}$ faithfully. This concludes the proof of Theorem 1.1. $\blacksquare$

We remark that Theorem 1.1 was already known when $\mathfrak{b}$ is compact (Gotay, Grabowski and Grundling [2000]), in which case the proof above simplifies greatly and provides an alternate means of establishing Theorem 2 *ibid.* Notice also that when $\mathfrak{b}$ is compact every quantization of $P(M)$ is necessarily polynomial; this follows from the observation that since $\mathcal{Q} \upharpoonright \mathfrak{b}$ is irreducible the representation space $\mathcal{H}$ must be finite-dimensional together with a well known fact about enveloping algebras (Prop. 2.6.5 in Dixmier [1976]).

3 Discussion

The key observation underlying Theorem 1.1 is that as $M \subset \mathfrak{b}^*$ is non-nilpotent, its ideal $I(M)$ is nonhomogeneous. If M is a basic nilpotent orbit, on the other hand, then $I(M)$ is homogeneous, and from Theorem 1.1 in Gotay [2002] we know that there do exist polynomial quantizations of $P(M)$. (Although it is not clear to what extent these are “nontrivial” in general.) Taken together, these two results serve to establish a conjecture of Gotay [2000] when $\mathfrak{b}$ is semisimple: *There exists a consistent polynomial quantization of $P(M)$ if and only if $I(M)$ is homogeneous.*

Finally, we remark that the restriction to basic coadjoint orbits is for physical reasons, cf. Gotay [2000]. For our purposes here, we may delete the adjective “basic” and consider arbitrary orbits. Both Theorem 1.1 and the conjecture above will remain valid in this extended context.

References

- Dixmier, J. [1977] *Enveloping Algebras*. (North Holland, Amsterdam).
- Gotay, M. J. [2000] Obstructions to quantization. In: *Mechanics: From Theory to Computation (Essays in Honor of Juan-Carlos Simo)*, J. Nonlinear Sci. Eds. (Springer, New York), 171–216.
- Gotay, M. J. [2002] On quantizing semisimple basic algebras, I: $\mathfrak{sl}(2, \mathbb{R})$. To appear in: *The Jerry Marsden 60th Birthday Volume*, P. Holmes, P. Newton, and A. Weinstein, Eds. (Springer, New York). Preprint math-ph/0012034.
- Gotay, M. J., Grabowski, J., and Grundling, H. B. [2000] An obstruction to quantizing compact symplectic manifolds. *Proc. Amer. Math. Soc.* **28**, 237–243.
- Varadarajan, V. S. [1984] *Lie Groups, Lie Algebras, and their Representations*. (Springer, New York).